

On the Simultaneous Distribution of the Fractional Parts of Different Powers of Prime Numbers¹

Wenguang Zhai

Department of Mathematics, Shandong Normal University, Jinan 250014, China

E-mail: wgzhai@jn-public.sd.cninfo.net

Communicated by A. Hildebrand

Received November 8, 1999

In this paper we study the distribution modulo 1 of the sequence of vectors



provided by Elsevier - Publisher Connector

1. INTRODUCTION AND MAIN RESULTS

In 1940 I. M. Vinogradov considered the distribution of the fractional parts of the sequence $f\sqrt{p}$, where p runs over prime numbers (see [15]). This celebrated work motivated the interests of many authors to investigate the distribution of p^θ modulo 1 by various methods (see [1–3, 7, 8, 10–12]).

In [13], Tolev studied the simultaneous distribution of the fractional parts of different powers of prime numbers. Let $k \geq 2$ be a fixed integer and $0 < \alpha_k < \dots < \alpha_1 < 0$ real numbers; $\Gamma \subset R^k$ is defined by

$$\Gamma = \Gamma(\xi_1, \eta_1, \dots, \xi_k, \eta_k) = \{(x_1, \dots, x_k) \mid \xi_i < x_i < \eta_i, 1 \leq i \leq k\}$$

where $0 < \xi_i < \eta_i \leq 1$, $1 \leq i \leq k$. Let $\mu(\Gamma) = \prod_{i=1}^k (\eta_i - \xi_i)$, and let $S(x; \Gamma)$ denote the number of primes not greater than x and satisfy the condition

$$(\{p^{\alpha_1}\}, \{p^{\alpha_2}\}, \dots, \{p^{\alpha_k}\}) \in \Gamma.$$

Then Tolev proved that

$$S(x; \Gamma) = \pi(x)(\mu(\Gamma) + O(x^{-\delta/3} \log^{k+9} x)) \quad (1.1)$$

¹ This work is supported by the National Natural Science Foundation of China (Grant 19801021) and the Natural Science Foundation of Shandong Province (Grant Q98A02110).



with

$$\delta = \min(1 - \alpha_1, \alpha_1 - \alpha_2, \dots, \alpha_{k-1} - \alpha_k, \alpha_k, 1/4).$$

The aim of this paper is to further improve Tolev's result by different methods. We first prove the following

THEOREM 1. *We have*

$$S(x; \Gamma) = \pi(x)(\mu(\Gamma) + O(x^{-\delta_1} \log^{k+11.5} x)), \quad (1.2)$$

where

$$\delta_1 = \min(1 - \alpha_1, \alpha_1 - \alpha_2, \dots, \alpha_{k-1} - \alpha_k, \alpha_k/3, 20/177).$$

EXAMPLE 1. Take $k = 2$. If $80/177 < \alpha_1 < 157/177$, $60/177 < \alpha_2 < \alpha_1 - 20/177$, then

$$S(x; \Gamma) = \pi(x) \mu(\Gamma) + O(x^{157/177} \log^{14.5} x). \quad (1.3)$$

Similar to Theorem 1, we can prove the following.

THEOREM 2. *We have*

$$S(x; \Gamma) = \pi(x)(\mu(\Gamma) + O(x^{-\delta_2} \log^{k+11.5} x)), \quad (1.4)$$

where

$$\delta_2 = \min(\alpha_1 - \alpha_2, \dots, \alpha_{k-1} - \alpha_k, \alpha_k/3, 40/407).$$

EXAMPLE 2. Take $k = 2$. If $120/407 < \alpha_2 < \alpha_1 - 40/407$, then

$$S(x; \Gamma) = \pi(x) \mu(\Gamma) + O(x^{367/407} \log^{14.5} x). \quad (1.5)$$

Obviously, Theorem 2 is better than Theorem 1 if α_1 is very close to 1.

It is obvious that Theorem 1 and Theorem 2 are very weak if $\delta_0 = \min(\alpha_1 - \alpha_2, \alpha_2 - \alpha_3, \dots, \alpha_{k-1} - \alpha_k)$ is very small. So we shall use a different approach to prove the following Theorem 3, which is better than Theorem 1 and Theorem 2 if δ_0 is small.

THEOREM 3. *We have*

$$S(x; \Gamma) = \pi(x)(\mu(\Gamma) + O(x^{-\delta_3} \log^{k+5.5} x)), \quad (1.6)$$

where

$$\delta_3 = \min(1/(4k+6), \alpha_k/(4k-2)).$$

EXAMPLE 3. Take $k = 2$. If $6/14 < \alpha_2 < \alpha_1 < 1$, $\alpha_1 - \alpha_2 < 1/14$. Then Theorem 3 yields

$$S(x; \Gamma) = \pi(x) \mu(\Gamma) + O(x^{13/14} \log^{8.5} x). \quad (1.7)$$

Notations. $\{l\}$ denotes the fractional part of t . $h = (h_1, \dots, h_k)$ denotes the k -dimensional vector with integer components.

$$\|h\| = \max_{1 \leq i \leq k} |h_i|; \quad r(h) = \prod_{i=1}^k \max(|h_i|, 1).$$

$\langle \cdot, \cdot \rangle$ denotes the Euclidean scalar product in R^k . $e(x) = e^{2\pi i x}$. $m \sim M$ means $M < m \leq 2M$; $m \asymp M$ means $c_1 M \leq m \leq c_2 M$ for positive constants c_1 and c_2 . $\Lambda(n)$ is the Mangoldt function.

2. SOME PRELIMINARY LEMMAS

We need the following lemmas.

LEMMA 2.1. Suppose $f(n)$ is a real-valued function on the interval $[N, N_1]$, where $2 \leq N < N_1 \leq 2N$. If $0 < c_1 \lambda_1 \leq |f'(n)| \leq c_2 \lambda_1 \leq 1/2$, then

$$\sum_{N < n \leq N_1} e(f(n)) \ll \lambda_1^{-1}.$$

If $|f^{(j)}(n)| \sim \lambda_1 N^{-j+1}$ ($j = 1, 2$), then

$$\sum_{N < n \leq N_1} e(f(n)) \ll \lambda_1^{-1} + N^{1/2} \lambda_1^{1/2}.$$

If $|f^{(j)}(n)| \sim \lambda_1 N^{-j+1}$ ($j = 1, 2, 3, 4, 5, 6$), then

$$\sum_{N < n \leq N_1} e(f(n)) \ll \lambda_1^{-1} + N^\lambda \lambda_1^\kappa.$$

Where (κ, λ) is any exponent pair.

Proof. See Graham and Kolesnik [6].

LEMMA 2.2. Suppose that $0 < a < b \leq 2a$ and R is an open convex set in C containing the real segment $[a, b]$. Suppose further that $f(z)$ is analytic on R , $f(x)$ is real for real x in R , $|f''(z)| \leq M$ for $z \in R$, and there is a constant

$k > 0$ such that $f''(x) \leq -kM$ for all real x in R . Let $\alpha = f'(b)$ and $\beta = f'(a)$, and define x_v for each integer v in the range $\alpha < v < \beta$ by $f'(x_v) = v$. Then we have

$$\sum_{a < n \leq b} e(f(n)) = e\left(-\frac{1}{8}\right) \sum_{\alpha < v \leq \beta} |f''(x_v)|^{-1/2} e(f(x_v) - vx_v) \\ + O(M^{-1/2} + \log(2 + M(b-a))).$$

Proof. This is Lemma 6 of Heath-Brown [9].

LEMMA 2.3. Suppose $z(n)$ is any complex number, $1 \leq Q \leq N$, then

$$\left| \sum_{N < n \leq CN} z(n) \right|^2 \ll \frac{N}{Q} \sum_{0 \leq q \leq Q} \left(1 - \frac{q}{Q}\right) \operatorname{Re} \sum_{N < n \leq CN-q} z(n) \overline{z(n+q)}.$$

Proof. This is Lemma 2.5 of Graham and Kolesnik [6].

LEMMA 2.4. Let $0 < L \leq N < vN \leq \lambda L$ and a_n be complex numbers with $|a_n| \leq 1$. Then we have

$$\sum_{N < n \leq vN} a_n = \frac{1}{2\pi} \int_{-L}^L \left(\sum_{L < l \leq L} a_l l^{-it} \right) \frac{(vN)^{it} - N^{it}}{t} dt + O(\log(2 + L)).$$

Proof. This is Lemma 6 of Fouvry and Iwaniec [5].

LEMMA 2.5. If $Z_n = (Z_{1,n}, \dots, Z_{k,n})$ ($n = 1, 2, 3, \dots$) is a sequence of k -dimensional vectors and

$$D_N = \sup_{\Gamma} \left| \frac{1}{N} \sum_{\substack{n \leq N \\ (Z_{1,n}, \dots, Z_{k,n}) \in \Gamma}} 1 - \mu(\Gamma) \right|$$

is its discrepancy (the supremum is taken over the set of all Γ of type (1.1)), then the inequality

$$D_N \ll \frac{1}{M} + \sum_{0 < \|m\| \leq M} \frac{1}{r(m)} \left| \frac{1}{N} \sum_{n \leq N} e(\langle m, Z_n \rangle) \right|$$

holds. (M is an arbitrary positive number; the constant implied in the \ll symbol depends only on k .)

Proof. This is Lemma 3 of Tolev [13].

LEMMA 2.6. *Let \mathcal{X} and \mathcal{Y} be two finite set of real numbers, $\mathcal{X} \subset [-X, X]$, and $\mathcal{Y} \subset [-Y, Y]$. Then for any complex function $u(x)$ and $v(y)$ we have*

$$\left| \sum_{x \in \mathcal{X}} \sum_{y \in \mathcal{Y}} u(x) v(y) e(xy) \right|^2 \leq 20(1 + XY) \sum_{\substack{x, x' \in \mathcal{X} \\ |x - x'| \leq Y^{-1}}} |u(x) u(x')| \sum_{\substack{y, y' \in \mathcal{Y} \\ |y - y'| \leq X^{-1}}} |v(y) v(y')|.$$

Proof. This is the one-dimensional case of Lemma 2.5 of [4]; see also Proposition 1 of Fouvry and Iwaniec [5].

LEMMA 2.7. *Let u and v be positive real numbers. If $n > v$ then*

$$\begin{aligned} \Lambda(n) = & - \sum_{\substack{n=kl \\ k > v, l > u}} \Lambda(k) \sum_{d|l, d \leq u} \mu(d) + \sum_{n=kl, l \leq u} \mu(k) \log l \\ & - \sum_{\substack{n=klm \\ l \leq u, m \leq v}} \Lambda(k) \mu(m). \end{aligned}$$

Proof. See Vaughan [14].

Suppose $d \geq 2$ is a fixed integer; a_1, a_2, \dots, a_d are any real numbers with $a_1 a_2 \cdots a_d \neq 0$; $\gamma_1, \dots, \gamma_d$ are real noninteger constants; and M and M_1 are real numbers such that $5 < M < M_1 \leq 2M$. Let

$$f_d(m) = a_1 m^{\gamma_1} + \cdots + a_d m^{\gamma_d}.$$

Set $R = |a_1| M^{\gamma_1} + \cdots + |a_d| M^{\gamma_d}$. For the exponential sum

$$S_d(m) = \sum_{M < m \leq M_1} e(f_d(m)),$$

we have the following two lemmas, which are Propositions 2 and 3 of Zhai [16] respectively.

LEMMA 2.8. *If $R \leq \eta M$, where η is a fixed positive constant small enough, then*

$$S_d(M) \ll MR^{-1/d}.$$

LEMMA 2.9. *We have*

$$S_d(M) \ll R^{1/2} + MR^{-1/(d+1)}.$$

3. ESTIMATION OF AN EXPONENTIAL SUM OVER PRIMES

Suppose Y is a large positive real number; $0 < \alpha < 1$; $0 < \delta(\alpha) < 1/3$ is a function of α ; h is an integer such that $1 \leq h \ll Y^\delta$; and $g(m)$ is a real function on $[Y, 2Y]$ of the form

$$g(m) = u_1 m^{\gamma_1} + \dots + u_l m^{\gamma_l}$$

such that $|g^{(j)}(m)| \leq \varepsilon h Y^{\alpha-j}$ ($j = 0, 1, 2, \dots, 6$), where $l \geq 1$, $\gamma_1, \dots, \gamma_l$ are real constants; u_1, \dots, u_l are any real numbers; and ε is a sufficiently small positive constant. The aim of this section is to estimate the exponential sum over primes

$$S(Y; h, \alpha) = \sum_{Y < m \leq 2Y} A(m) e(hm^\alpha + g(m)).$$

For convenience, in this section we always set $F = hY^\alpha$. The constants implied by \ll in this section depend only on $\alpha, \gamma_1, \dots, \gamma_l$ and ε .

LEMMA 3.1. *Suppose $340/531 < \alpha < 1$, $\delta = \delta(\alpha) = \min(1 - \alpha, 20/177)$, and $0 < A \leq \delta$. Let a_m be a sequence of complex numbers such that*

$$\sum_{m \sim M} |a_m|^2 \ll M \log^{2A} M, \quad A > 0.$$

Then for $h \ll Y^\delta$, $M \ll Y^{1-2A}$, we have

$$S_I = \sum_{m \sim M} a(m) \sum_{Y < mn \leq 2Y} e(h(mn)^\alpha + g(mn)) \ll Y^{1-A} \log^{A+1} Y. \quad (3.1)$$

Proof. Write $f(m) = hm^\alpha + g(m)$ and let $N = Y/M$.

By Lemma 2.1 with the exponent pair $(2/7, 4/7)$ we get

$$\begin{aligned} S_I &\ll \sum_{m \sim M} |a_m| (NF^1 + F^{2/7} N^{2/7}) \\ &\ll MNF^{-1} \log^A Y + MF^{2/7} N^{2/7} \log^A Y \\ &\ll Y^{1-\alpha} \log^A Y + Y^{1-A} \log^A Y \ll Y^{1-A} \log^A Y, \end{aligned} \quad (3.2)$$

if $M \ll Y^{(5-7A)/5} F^{-2/5}$.

Now suppose $M \gg Y^{(5-7A)/5} F^{-2/5}$. Take $Q = [Y^{2A} \log^{-1} Y]$, then $Q = o(N)$. By Cauchy's inequality and Lemma 2.3 we get

$$\begin{aligned} |S_I|^2 &\ll \sum_{m \sim M} |a_m|^2 \sum_{m \sim M} \left| \sum_{n \sim Y/m} e(f(mn)) \right|^2 \\ &\ll \frac{M^2 N^2 \log^{2A} M}{Q} + \frac{MN \log^{2A} M}{Q} \sum_{q=1}^Q |E_q|, \end{aligned} \quad (3.3)$$

where

$$\begin{aligned} E_q &= \sum_{M < m \leq 2M} \sum_{Y/m < n \leq 2Y/m - q} e \left(hm^\alpha \Delta(n, q; \alpha) + \sum_{j=1}^l u_j m^{\gamma_j} \Delta(n, q; \gamma_j) \right), \\ \Delta(n, q; t) &= (n+q)^t - n^t. \end{aligned}$$

So the problem is reduced to show that

$$\sum_{q=1}^Q |E_q| \ll Y. \quad (3.4)$$

For fixed q , write

$$G(m, n) = G(m, n; q) = hm^\alpha \Delta(n, q; \alpha) + \sum_{j=1}^l u_j m^{\gamma_j} \Delta(n, q; \gamma_j).$$

If $\max |\partial G / \partial m| \leq 1/2$, then by Lemma 2.1 we get

$$E_q \ll \frac{MN^2}{Fq}. \quad (3.5)$$

If $|\partial G / \partial n| \leq 1/2$, we can get the same estimate.

Now suppose $|\partial G / \partial m| \gg 1$, $|\partial G / \partial n| \gg 1$. We have

$$E_q = \sum_{Y/2M < n \leq 2Y/M - q} \sum_{M_1 < m \leq M_2} e(G(m, n)),$$

where $M_1 = \max(M, Y/n)$, $M_2 = \min(2M, 2Y/(n+q))$. Using Lemma 2.2 we have

$$\begin{aligned} &\sum_{M_1 < m \leq M_2} e(G(m, n)) \\ &= c \sum_{r_1(n) < r \leq r_2(n)} \left| \frac{\partial^2 G}{\partial m^2}(m_r, n) \right|^{-1/2} e(G(m_r, n) - rm_r) \\ &\quad + O(\log Y + MN^{1/2} (Fq)^{-1/2}), \end{aligned}$$

where $m_r = m(r, n)$ is the solution of the equation $(\partial G / \partial m)(m, n) = r$ and

$$r_1(n) = \frac{\partial G}{\partial m}(M_1, n), \quad r_2(n) = \frac{\partial G}{\partial m}(M_2, n).$$

Let $r_1 = \min_n r_1(n)$, $r_2 = \min_n r_2(n)$, and

$$\varphi(n, r) = \frac{(Fq)^{1/2}}{MN^{1/2}} \left| \frac{\partial^2 G}{\partial m^2}(m_r, n) \right|^{-1/2}, \quad s(r, n) = G(m_r, n) - rm_r.$$

For each $r_1 \leq r \leq r_2$, the equations

$$Y/2M < n \leq 2Y/M - q, \quad r_1(n) < r \leq r_2(n)$$

define an interval $I(r) = (n_1(r), n_2(r)]$.

So we have

$$\begin{aligned} E_q &\ll \frac{MN^{1/2}}{(Fq)^{1/2}} \sum_{r_1 \leq r \leq r_2} \left| \sum_{n \in I(r)} \varphi(n, r) e(s(r, n)) \right| \\ &\quad + N \log Y + MN^{3/2} (Fq)^{-1/2}, \end{aligned}$$

where $I(r) = (n_1(r), n_2(r)]$.

We can verify that

$$\begin{aligned} \varphi(n, r) &\ll 1, \quad \varphi'(n, r) \ll N^{-1}, \\ \frac{\partial^j s}{\partial n^j}(r, n) &\sim FqN^{-j-1}, \quad j = 0, 1, 2, 3, 4, 5, 6. \end{aligned}$$

So by Lemma 2.1 with the exponent $(2/18, 13/18)$ we get

$$\begin{aligned} E_q &\ll \frac{MN^{1/2}}{(Fq)^{1/2}} \frac{Fq}{MN} \left(\frac{Fq}{N^2} \right)^{2/18} N^{13/18} \\ &\quad + N \log Y + MN^{3/2} (Fq)^{-1/2} \\ &\ll (Fq)^{11/18} + N \log Y + MN^{3/2} (Fq)^{-1/2}. \end{aligned} \tag{3.6}$$

Combining (3.5) and (3.6) we have

$$\begin{aligned} \sum_{q=1}^Q |E_q| &\ll \frac{MN^2 \log Q}{F} + F^{11/18} Q^{29/18} + NQ \log Y \\ &\quad + MN^{3/2} Q^{1/2} (F)^{-1/2} = \sum_{j=1}^4 I_j. \end{aligned}$$

Since $\alpha \geq 340/531$, we have $\Delta \leq \delta \leq 3\alpha/17$, which gives $M \gg Y^{(5-7\Delta)/5} F^{-2/5} \gg YQ/F$. So we have $I_1 \ll Y$, $I_4 \ll Y$. Since $11\alpha + 69\delta \leq 18$, we have $I_2 \ll Y$. Since $2\alpha + 19\delta < 5$, we have $M \gg Y^{(5-7\Delta)/5} F^{-2/5} \gg Q \log Y$, so $I_3 \ll Y$. Combining the above, we get (3.4). This completes the proof of Lemma 3.1.

LEMMA 3.2. Suppose $340/531 < \alpha < 1$, $\delta = \delta(\alpha) = \min(1 - \alpha, 20/177)$, $0 < \Delta \leq \delta$. Let $a(m)$ and $b(n)$ be complex numbers such that

$$\sum_{M < m \leq 2M} |a(m)|^2 \ll M \log^{2A} M,$$

$$\sum_{N < n \leq 2N} |b(n)|^4 \ll N \log^{4B} N, \quad A > 0, \quad B > 0.$$

Then for $Y^{2\Delta} \ll N \ll \min(FY^{-2\Delta}, Y^{1-4\Delta}, Y^{(93-318\Delta)/9} F^{-53/9})$, $MN \sim Y$, we have

$$S_{\Pi} = \sum_{m \sim M} a(m) \sum_{n \sim N} b(n) e(h(mn)^\alpha + g(mn))$$

$$\ll Y^{1-\Delta} \log^{A+B+1} Y. \quad (3.7)$$

Proof. Take $Q = [Y^{2\Delta} \log^{-1} Y]$, then $Q = o(N)$. By Cauchy's inequality and Lemma 2.3 we get

$$|S_{\Pi}|^2 \ll \sum_{m \sim M} |a(m)|^2 \sum_{m \sim M} \left| \sum_{n \sim N} b(n) e(f(mn)) \right|^2$$

$$\ll \frac{M^2 N^2 \log^{2A+2B} Y}{Q} + \frac{MN \log^{2A} Y}{Q} \sum_{q=1}^Q E_q, \quad (3.8)$$

where

$$E_q = \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{m \sim M} e(G(m, n)) \right|$$

and $G(m, n)$ is defined in the same way as in the proof of Lemma 3.1.

If $|\partial G / \partial m| \leq 10^3 M q^{-2}$, by Lemma 2.1 we get

$$E_q \ll \sum_{n \sim N} |b(n+q) b(n)| \left(\frac{MN}{Fq} + \left(\frac{Fq}{MN} \right)^{1/2} M^{1/2} \right)$$

$$\ll \sum_{n \sim N} (|b(n+q)|^2 + |b(n)|^2) \left(\frac{MN}{Fq} + M/q \right)$$

$$\ll MN/q \log^{2B} Y, \quad (3.9)$$

if we note that $N \ll F$.

Now suppose $|\partial G/\partial m| > 10^3 Mq^{-2}$. By Lemma 2.2 we get

$$\sum_{m \sim M} e(G(m, n)) \ll \frac{MN^{1/2}}{(Fq)^{1/2}} \left| \sum_{r_1(n) \leq r \leq r_2(n)} \varphi(n, r) e(s(r, n)) \right| + \log Y + MN^{1/2} (Fq)^{-1/2},$$

where $s(r, n)$, $\varphi(n, r)$ are defined in the same way as in the proof of Lemma 3.1 and

$$r_1(n) = \frac{\partial G}{\partial m}(M, n), \quad r_2(n) = \frac{\partial G}{\partial m}(2M, n).$$

Thus we have

$$E_q \ll \frac{MN^{1/2}}{(Fq)^{1/2}} \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{r_1(n) < r \leq r_2(n)} \varphi(n, r) e(s(r, n)) \right| + N \log^{2B+1} Y + MN^{3/2} (Fq)^{-1/2} \log^{2B} Y. \quad (3.10)$$

So it suffices to bound the sum

$$\Sigma_1 = \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{r_1(n) < r \leq r_2(n)} \varphi(n, r) e(s(r, n)) \right|.$$

By Cauchy's inequality and Lemma 2.3 again we get

$$\begin{aligned} \Sigma_1^2 &\ll \sum_{n \sim N} |b(n+q) b(n)|^2 \sum_{n \sim N} \left| \sum_{r_1(n) < r \leq r_2(n)} \varphi(n, r) e(s(r, n)) \right|^2 \\ &\ll \frac{N^2 R^2 \log^{4B} Y}{T} + \frac{NR \log^{4B} Y}{T} \Sigma_2 \end{aligned} \quad (3.11)$$

with $T = [Fq^3/M^2N]$ and $R = Fq/MN$, where

$$\Sigma_2 = \sum_{t=1}^T \left| \sum_{n \sim N} \sum_{r_1(n) < r \leq r_2(n)-t} \varphi(n, r+t) \varphi(n, r) e(s(r+t, n) - s(r, n)) \right|$$

and where we used the estimate

$$\sum_{n \sim N} |b(n+q) b(n)|^2 \ll \sum_{n \sim N} (|b(n+q)|^4 + |b(n)|^4) \ll N \log^{4B} Y.$$

It is easy to check that $10 < T = o(R)$.

We recall that $s(r, n) = G(m(r, n), n) - rm(r, n)$, where $m(r, n)$ denotes the solution of

$$\frac{\partial G}{\partial m}(m, n) = r.$$

It can be easily seen that

$$\frac{\partial s}{\partial r}(r, n) = \frac{\partial G}{\partial m} \frac{\partial m}{\partial r} - m(r, n) - r \frac{\partial m}{\partial r} = -m(r, n).$$

So we have

$$\begin{aligned} H(n) &:= H_{r,t,q}(n) = s(r+t, n) - s(r, n) \\ &= \int_r^{r+t} \frac{\partial s}{\partial u}(u, n) du = - \int_r^{r+t} m(u, n) du, \end{aligned}$$

which implies that

$$|H^{(j)}(n)| \sim tMN^{-j} \quad (j=0, 1, 2, 3, 4, 5, 6.)$$

Let $I(r, t)$ denote the interval

$$N < n \leq 2N, \quad r_1(n) < r \leq r_2(n) - t.$$

Then

$$\Sigma_2 \ll \sum_{t=1}^T \sum_{r \sim R} \left| \sum_{n \in I(r, t)} \varphi(n, r+t) \varphi(n, r) e(s(r+t, n) - s(r, n)) \right|.$$

Thus using partial summation and then using the exponent pair $(13/40, 22/40) = (BA^2)^2(1/2, 1/2)$, we get

$$\begin{aligned} \Sigma_2 &\ll \sum_{t=1}^T \sum_{r \sim R} (tM/N)^{13/40} N^{22/40} \\ &\ll RM^{13/40} N^{9/40} T^{53/40} \ll NR \end{aligned} \quad (3.12)$$

if we note that $N \ll Y^{(93-318A)/9} F^{-53/9}$.

Combining (3.9)–(3.12) we get that for any $1 \leq q \leq Q$,

$$\begin{aligned} E_q &\ll \frac{MN \log^{2B+1} Y}{q} + N \log^{2B+1} Y \\ &\quad + MN^{3/2} (Fq)^{-1/2} \log^{2B} Y. \end{aligned} \quad (3.13)$$

Now Lemma 3.2 follows from inserting (3.13) into (3.8).

From Lemma 3.1 and Lemma 3.2 we can prove the following Proposition 3.1.

PROPOSITION 3.1. *Suppose $340/531 < \alpha < 1$, $\delta = \delta(\alpha) = \min(1 - \alpha, 20/177)$, $0 < \Delta \leq \delta$. Then, for $h \ll Y^\delta$, we have*

$$S(Y; h, \alpha) \ll Y^{1-\Delta} \log^{11.5} Y. \quad (3.14)$$

Proof. Applying Lemma 2.7 to $S(Y, h, \alpha)$ we get

$$\begin{aligned} S(Y, h, \alpha) &= \sum_{m \leq u} \sum_{n \sim Y/m} \mu(m) \log ne(f(mn)) \\ &\quad - \sum_{m \leq uv} \sum_{n \sim Y/m} b(m) e(f(mn)) \\ &\quad - \sum_{u < n \leq 2Y/v} \sum_{m \sim Y/n, m > v} a(n) \Lambda(m) e(f(mn)) \\ &= S_1 - S_2 - S_3, \end{aligned} \quad (3.15)$$

where

$$a(m) = \sum_{d|m, d \leq u} \mu(d), \quad b(n) = \sum_{n=de, d \leq u, e \leq v} \mu(d) \Lambda(e).$$

Note that $|a(m)| \leq d(m)$ and $|b(m)| \leq \sum_{d|m} \Lambda(d) = \log m$. So, for any $D \geq 2$,

$$\sum_{m \sim D} |a(m)|^2 \ll D \log^3 D, \quad \sum_{n \sim D} |b(n)|^2 \ll D \log^2 D.$$

It suffices to bound S_i ($i = 1, 2, 3$). We take $u = Y^{2\Delta}$, $v = Y/E$, where

$$E = \min(FY^{-2\Delta}, Y^{1-4\Delta}, Y^{(93-318\Delta)/9} F^{-53/9}).$$

Using Lemma 1 with the exponent pair $(1/2, 1/2)$ we get

$$\begin{aligned} S_1 &\ll \sum (Y/mF + F^{1/2}) \log Y \\ &\ll (Y/F \log^2 Y + uF^{1/2} \log Y) \ll Y^{1-\Delta} \log^2 Y. \end{aligned}$$

S_2 can be divided into $O(\log Y)$ sums of the form

$$S_2(M) = \sum_{m \sim M} b(m) \sum_{n \sim Y/m} e(f(mn)) \quad (M \ll uv). \quad (3.16)$$

It is easy to show that $uv \ll Y^{1-2A}$. So by Lemma 3.1 each subsum is $\ll Y^{1-A} \log^2 Y$. After adding these subsums, we find that

$$S_2 \ll Y^{1-A} \log^3 Y.$$

S_3 can be divided into $O(\log Y)$ sums of the form

$$S_3(N) = \sum_{n \sim N} \sum_{m \sim Y/n, m > v} a(n) A(m) e(f(mn))$$

$$(u \ll N \ll Y/v). \quad (3.17)$$

By Lemma 2.4

$$\sum_{m \sim Y/n, m > v} A(m) e(f(mn))$$

$$= \frac{1}{2\pi} \int_{-M}^M \left(\sum_{M < m < 4M} A(m) m^{it} e(f(mn)) \right)$$

$$\times \frac{m_2^{it}(n) - m_1^{it}(n)}{t} dt + O(\log^2 Y)$$

with $M = Y/2N$.

So we have

$$S_3(N) = \frac{1}{2\pi} \int_{-M}^M \sum_{n \sim N} a(n) \sum_{M < m < 4M} A(m) m^{it} e(f(mn))$$

$$\times \frac{m_2^{it}(n) - m_1^{it}(n)}{t} dt + O(\log^2 Y)$$

$$\ll |S_3^*(N)| \log Y + HN \log^4 Y, \quad (3.18)$$

where

$$S_3^*(N) = \sum_{n \sim N} \sum_{m \sim M} a^*(n) b^*(m) e(f(mn)), \quad (3.19)$$

with $a^*(n) \ll d(n)$, $b^*(m) \ll \log m$. So we have

$$\sum_{n \sim N} |a^*(n)|^4 \ll N \log^{15} N, \quad \sum_{m \sim M} |b^*(m)|^2 \ll M \log^2 M.$$

By Lemma 3.2 we get $S_3^*(N) \ll Y^{1-A} \log^{9.5} Y$. Combining the above we get

$$S_3 \ll Y^{1-A} \log^{11.5} Y.$$

This completes the proof of Proposition 3.1.

By the same proof we can prove the following.

PROPOSITION 3.1*. *Suppose $340/531 < \alpha < 1$, $\delta = 40/407$, $0 < \Delta \leq \delta$. Then for $h \ll Y^\delta$, we have*

$$S(Y; h, \alpha) \ll Y^{1-\Delta} \log^{11.5} Y.$$

LEMMA 3.3. *Suppose $0 < \alpha < 4/5$, $\delta = \delta(\alpha) = \min((1-\alpha)/3, \alpha/4)$, $0 < \Delta \leq \delta$. Let a_m be a sequence of complex numbers such that*

$$\sum_{M < m \leq 2M} |a_m|^2 \ll M \log^{2A} X, \quad A > 0.$$

Then for $h \ll Y^\delta$, $M \ll YF^{-1}$, we have

$$S_I = \sum_{m \sim M} a(m) \sum_{n \sim Y/m} e(h(mn)^\alpha + g(mn)) \ll Y^{1-\Delta} \log^4 Y. \quad (3.20)$$

Proof. It immediately follows from Lemma 2.1.

LEMMA 3.4. *Suppose $0 < \alpha < 4/5$, $\delta = \delta(\alpha) = \min((1-\alpha)/3, \alpha/4)$, $0 < \Delta \leq \delta$. Let $a(m)$ and $b(n)$ be complex numbers such that*

$$\begin{aligned} \sum_{M < m \leq 2M} |a(m)|^2 &\ll M \log^{2A} M, \\ \sum_{N < n \leq 2N} |b(n)|^2 &\ll N \log^{2B} N, \quad A > 0, \quad B > 0. \end{aligned}$$

Then for $h \ll Y^\delta$, $Y^{2\Delta} \ll N \ll F$, we have

$$\begin{aligned} S_{II} &= \sum_{m \sim M} a(m) \sum_{n \sim N} b(n) e(h(mn)^\alpha + g(mn)) \\ &\ll Y^{1-\Delta} \log^{A+B+1} Y. \end{aligned} \quad (3.21)$$

Proof. We begin with (3.8). Since $\Delta \leq (1-\alpha)/3$, we have $|\partial G / \partial m| \leq 1/2$. By Lemma 2.1 we have

$$\begin{aligned} \sum_{q=1}^Q E_q &\ll \sum_{q=1}^Q \sum_{n \sim N} |b(n+q) b(n)| \frac{MN}{Fq} \\ &\ll MN^2 F^{-1} \log^{2B+1} Y \ll Y \log^{2B+1} Y. \end{aligned} \quad (3.22)$$

Whence Lemma 3.4 follows.

PROPOSITION 3.2. *Suppose $0 < \alpha < 4/5$, $\delta = \min((1 - \alpha)/3, \alpha/4)$, $0 < \Delta \leq \delta$. Then for $h \ll Y^\delta$ we have*

$$S(Y, h, \alpha) \ll Y^{1-\Delta} \log^{5.5} Y. \quad (3.23)$$

Proof. We use notations in the proof of Proposition 3.1. Take $u = Y^{2\Delta}$, $v = Y/F$.

By Lemma 2.1 we have

$$S_1 \ll Y^{1-\Delta} \log Y.$$

To estimate S_2 , we only need to estimate $S_2(M)$. If $M \ll Y/F = v$, by Lemma 3.3 we get $S_2(M) \ll Y^{1-\Delta} \log^2 Y$. If $v \ll M \ll uv$, then $Y^{2\Delta} \ll N = Y/M \ll F$. So by Lemma 3.4 we can get the same estimate. Thus we have

$$S_2 \ll Y^{1-\Delta} \log^3 Y.$$

By Lemma 3.4 we have

$$S_3^*(N) \ll Y^{1-\Delta} \log^{3.5} Y.$$

Thus

$$S_3 \ll Y^{1-\Delta} \log^{5.5} Y.$$

This completes the proof of Proposition 3.2.

PROPOSITION 3.3. *Suppose $0 < \alpha < 2/3$, $\delta = \min((1 - \alpha)/3, \alpha/2, 1/6)$. Then for $h \ll Y^\delta$, we have*

$$\sum_{m \sim M} \Lambda(m) e(hm^\alpha) \ll Y^{1-\delta} \log^{4/5} Y. \quad (3.24)$$

Proof. We begin with (3.15) by choosing $u = v = Y^{1/3}$.

By Lemma 2.1 with the exponent pair $(1/2, 1/2)$ we have

$$\begin{aligned} S_1 &\ll \sum_{m \leq Y^{1/3}} (Y/mF + F^{1/2}) \ll YF \log F + Y^{1/3} F^{1/2} \\ &\ll Y^{1-\alpha} \log Y + Y^{7/9} \ll Y^{1-\delta} \log Y. \end{aligned}$$

Now we estimate $S_3^*(N)$ for $Y^{1/3} \ll N \ll Y^{2/3}$. By Lemma 2.6 we have

$$|S_3^*(N)|^2 \ll F \mathcal{A} \mathcal{B},$$

where

$$\begin{aligned} \mathcal{A} &= \sum_{|n_1^\alpha - n_1^\alpha| \leq 1/hM^\alpha} |a^*(n_1) a^*(n_2)| \ll \sum_{|n_1 - n_2| \ll N/F} d^2(n_1) \\ &\ll \sum_{n_1 \sim N} d^2(n_1) \sum_{|n_2 - n_1| \ll N/F} 1 \ll N(1 + N/F) \log^3 N, \\ \mathcal{B} &\ll \sum_{|m_1^\alpha - m_1^\alpha| \leq 1/hN^\alpha} |b(m_1) b(m_2)| \ll M(1 + M/F) \log^2 Y. \end{aligned}$$

So we get

$$\begin{aligned} S^*(N) \log^{-5/2} Y &\ll (FMN)^{1/2} + MN^{1/2} + M^{1/2}N + MN/F^{1/2} \\ &\ll Y^{1-\delta}, \end{aligned}$$

which implies

$$S_3 \ll Y^{1-\delta} \log^{4.5} Y.$$

Finally, we estimate S_2 . If $M \ll Y^{1/3}$, then similar to S_1 we have $S_2(M) \ll Y^{1-\alpha} \log Y + Y^{7/9} \log Y$. If $Y^{1/3} \ll M$, then similar to $S_3^*(N)$ we have $S_2(M) \ll Y^{1-\delta} \log^2 Y$. Thus we get

$$S_2 \ll Y^{1-\delta} \log^3 Y.$$

This completes the proof of Proposition 3.3.

4. PROOFS OF THEOREM 1 AND THEOREM 2

We only prove Theorem 1. It suffices to establish the inequality

$$R(Y) \ll Y^{-\delta_1} \log^{k+11.5} Y. \quad (4.1)$$

for all $Y \in [x^{1-\delta_1}, x]$, where

$$R(Y) = \sup_r \left| \frac{1}{\pi(2Y) - \pi(Y)} (S(2Y; \Gamma) - S(Y; \Gamma)) - \mu(\Gamma) \right|.$$

According to Lemma 2.5 we have

$$\begin{aligned}
 R(Y) &\ll H^{-1} + \sum_{0 < \|h\| \leq H} \frac{1}{r(h)} \\
 &\quad \times \left| \frac{1}{\pi(2Y) - \pi(Y)} \sum_{Y < p \leq 2Y} e(h_1 p^{\alpha_1} + \dots + h_k p^{\alpha_k}) \right| \\
 &\ll H^{-1} + Y^{-1/2} \log^{k+2} Y + Y^{-1} \log Y \sum_{0 < \|h\| \leq H} \frac{1}{r(h)} |U(h)| \quad (4.2)
 \end{aligned}$$

for every $H > 2$, where

$$\begin{aligned}
 U(h) &= \sum_{Y < n \leq 2Y} \Lambda(n) e(H(n)), \\
 H(t) &= h_1 t^{\alpha_1} + \dots + h_k t^{\alpha_k}.
 \end{aligned}$$

Propositions from the past section can be applied to estimate $U(h)$. We take $H = CY^{\delta_1}$, where C is a sufficiently large positive constant.

Let $h = (h_1, \dots, h_k)$ satisfy $0 < \|h\| \leq H$ and d be the first integer j with $h_j \neq 0$, then $H(t) = h_d t^{\alpha_d} + g(t)$. Since $\delta_1 \leq \alpha_d - \alpha_{d+1}$, we have $g(t) = O(C^{-1} |h_d| Y^{\alpha_d})$.

If $\alpha_d > 340/531$, we use Proposition 3.1 to estimate $U(h)$. We take $\Delta = \alpha_d - \alpha_{d+1}$ if $\alpha_d - \alpha_{d+1} \leq \min(1 - \alpha_d, 20/177)$, and $\Delta = \min(1 - \alpha_d, 20/177)$ otherwise. We get

$$\begin{aligned}
 U(h) &\ll Y^{1 - \min(1 - \alpha_d, \alpha_d - \alpha_{d+1}, 20/177)} \log^{11.5} Y \\
 &\ll Y^{1 - \min(1 - \alpha_1, \alpha_d - \alpha_{d+1}, 20/177)} \log^{11.5} Y \\
 &\ll Y^{1 - \delta_1} \log^{11.5} Y.
 \end{aligned}$$

Now suppose $\alpha_d \leq 340/531$. If $h_{d+1} = \dots = h_k = 0$, then by Proposition 3.3 we get

$$\begin{aligned}
 U(h) &\ll Y^{1 - \min((1 - \alpha_d)/3, 1/6, \alpha_d/2)} \log^{4.5} Y \\
 &\ll Y^{1 - \min(\alpha_k/2, 191/1593)} \log^{4.5} Y \\
 &\ll Y^{1 - \delta_1} \log^{4.5} Y.
 \end{aligned}$$

If there is at least one $h_j \neq 0$ ($j > d$), then $d \leq k - 1$. By Proposition 3.2 we have

$$U(h) \ll Y^{1 - \min((1 - \alpha_d)/3, \alpha_d - \alpha_{d+1}, \alpha_d/4)} \log^{5.5} Y.$$

If $\alpha_d - \alpha_{d+1} \leq \alpha_d/4$, then

$$\min((1 - \alpha_d)/3, \alpha_d - \alpha_{d+1}, \alpha_d/4) = \min((1 - \alpha_d)/3, \alpha_d - \alpha_{d+1}).$$

If $\alpha_d - \alpha_{d+1} > \alpha_d/4$, then

$$\alpha_d/4 \geq \alpha_{d+1}/3 \geq \alpha_k/3.$$

So we have

$$\begin{aligned} U(h) &\ll Y^{1 - \min((1 - \alpha_d)/3, \alpha_d - \alpha_{d+1}, \alpha_d/4)} \log^{5.5} Y \\ &\ll Y^{1 - \min((1 - \alpha_d)/3, \alpha_d - \alpha_{d+1}, \alpha_k/3)} \log^{5.5} Y \\ &\ll Y^{1 - \delta_1} \log^{5.5} Y. \end{aligned}$$

This completes the proof of Theorem 1.

Using Proposition 3.1* in the proof of Theorem 1 we get Theorem 2.

5. EXPONENTIAL SUMS OVER PRIMES

Suppose $l \geq 2$ is a fixed integer; $1 > \gamma_1 > \gamma_2 > \dots > \gamma_l > 0$ are real numbers; Y is a large positive number; and $0 < \delta = \delta(\gamma_1) < 1/2$ is a constant depending only on γ_1 . The aim of this section is to estimate the exponential sum

$$S(Y; h_1, \dots, h_l, \gamma_1, \dots, \gamma_l) = \sum_{Y < n \leq 2Y} A(n) e\left(\sum_{j=1}^l h_j n^{\gamma_j}\right),$$

where h_j are real numbers such that $1 \leq |h_j| \leq Y^{\delta(\gamma_1)}$, $j = 1, 2, \dots, l$. In this section we always set $R = \sum_{j=1}^l |h_j| Y^{\gamma_j}$.

LEMMA 5.1. *Let $\delta = \delta(\gamma_1) = \min(\gamma_1/(4l-2), (1-\gamma_1)/3)$ and a_m be a sequence of complex numbers such that*

$$\sum_{M < m \leq 2M} |a_m|^2 \ll M \log^{2A} M, \quad A > 0.$$

Then for $M \ll Y^{1-\delta} R^{-1/2}$, $MN = Y$, we have

$$S_I = \sum_{m \sim M} a(m) \sum_{n \sim Y/m} e\left(\sum_{j=1}^l h_j (mn)^{\gamma_j}\right) \ll Y^{1-\delta} \log^A Y. \quad (5.1)$$

Proof. By Lemma 2.9 we have

$$\begin{aligned} S_I &\ll \sum_{M < m \leq 2M} |a_m| (R^{1/2} + NR^{-1/(k+1)}) \\ &\ll MR^{1/2} \log^A Y + Y^{1-(\gamma_1/(k+1))} \log^A Y. \end{aligned}$$

Whence Lemma 5.1 follows.

LEMMA 5.2. Let $\delta = \delta(\gamma_1) = \min(\gamma_1/(4l-2), (1-\gamma_1)/3)$ and $a(m)$ and $b(n)$ be complex numbers such that

$$\begin{aligned} \sum_{M < m \leq 2M} |a(m)|^2 &\ll M \log^{2A} M, \\ \sum_{N < n \leq 2N} |b(n)|^2 &\ll N \log^{2B} N, \quad A > 0, \quad B > 0. \end{aligned}$$

Then for $Y^{2\delta} \ll N \ll RY^{-2(l-1)\delta}$, $MN = Y$, we have

$$\begin{aligned} S_{II} &= \sum_{m \sim M} a(m) \sum_{n \sim N} b(n) e\left(\sum_{j=1}^l h_j(mn)^{\gamma_j}\right) \\ &\ll Y^{1-\delta} \log^{A+B+1} Y. \end{aligned} \quad (5.2)$$

Proof. Take $Q = [Y^{2\delta} \log^{-1} Y]$, then $10 < Q = o(N)$. By Cauchy's inequality and Lemma 2.3 we get

$$|S_{II}|^2 \ll \frac{M^2 N^2 \log^{2A+2B} Y}{Q} + \frac{MN \log^{2A} Y}{Q} \sum_{q=1}^Q E_q, \quad (5.3)$$

with

$$E_q = \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{m \sim M} e(f(m, n, q)) \right|,$$

where

$$f(m, n, q) = \sum_{j=1}^l h_j m^{\gamma_j} \Delta(n, q; \gamma_j)$$

and $\Delta(n, q; \gamma_j)$ is defined in the same way as in Section 3.

For each fixed q , we find that

$$\begin{aligned}
 & \sum_{j=1}^l |h_j m^{\gamma_j} \Delta(n, q; \gamma_j)| \\
 &= \sum_{j=1}^l |h_j| m^{\gamma_j} (\gamma_j q n^{\gamma_j-1} + O(q^2 N^{\gamma_j-2})) \\
 &\asymp \frac{Rq}{N} = R^*(q),
 \end{aligned}$$

say. Since

$$\frac{R^*(q)}{N} \ll \frac{Rq}{MN} \ll \frac{Y^{\delta+\gamma_1} q}{MN} \ll \frac{Y^{\delta+\gamma_1+2\delta}}{MN \log Y} \ll \log^{-1} Y,$$

by Lemma 2.8 we get

$$\begin{aligned}
 & \sum_{q=1}^Q \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{m \sim M} e(f(m, n, q)) \right| \\
 &\ll \sum_{q=1}^Q \sum_{n \sim N} |b(n+q) b(n)| M(R^*(q))^{-1/l} \\
 &\ll MN^{1+1/l} Q^{1-1/l} R^{-1/l} \log^{2B} Y \ll Y \log^{2B} Y,
 \end{aligned} \tag{5.4}$$

where in the last step we used the assumption that $N \ll RY^{-2(l-1)\delta}$. So Lemma 5.2 follows from (5.3) and (5.4).

LEMMA 5.3. *The estimate*

$$S(Y; h_1, \dots, h_l, \gamma_1, \dots, \gamma_l) \ll Y^{1-\min((1-\gamma_1)/3, \gamma_1/(4l-2))} \log^{5.5} Y. \tag{5.5}$$

Proof. We only sketch the proof. We begin with (3.15) by choosing $u = Y^{2\delta}$, $v = Y^{1+2(l-1)\delta} R^{-1}$. We use Lemma 2.8 to estimate S_1 and Lemma 5.2 to estimate S_3 . We use Lemma 5.1 to estimate S_2 for $M \ll Y^{1-\delta} R^{-1/2}$ and Lemma 5.2 to estimate S_2 for $Y^{1-\delta} R^{-1/2} \ll M \ll uv$.

LEMMA 5.4. *Suppose $1/2 < \gamma_1 < 1$, $\delta = \delta(\gamma_1) = \min((3-2\gamma_1)/14, (\gamma_1-1/2)/2l)$. Let $a(m)$ and $b(n)$ be complex numbers such that*

$$\begin{aligned}
 & \sum_{M < m \leq 2M} |a(m)|^2 \ll M \log^{2A} M, \\
 & \sum_{N < m \leq 2N} |b(n)|^2 \ll N \log^{2B} N, \quad A > 0, \quad B > 0.
 \end{aligned}$$

Then for $Y^{2\delta} \ll N \ll Y^{1/2}$, $MN \sim Y$, we have

$$\begin{aligned} S_{\text{II}} &= \sum_{m \sim M} a(m) \sum_{n \sim N} b(n) e \left(\sum_{j=1}^l h_j (mn)^{\gamma_j} \right) \\ &\ll Y^{1-\delta} \log^{A+B+1} Y. \end{aligned} \quad (5.6)$$

Proof. Using Lemma 2.7 to estimate the right side of (5.3) we get

$$\begin{aligned} &\sum_{q=1}^Q \sum_{n \sim N} |b(n+q) b(n)| \left| \sum_{m \sim M} e(f(m, n, q)) \right| \\ &\ll \sum_{q=1}^Q \sum_{n \sim N} |b(n+q) b(n)| \\ &\quad \times ((Rq)^{1/2} N^{-1/2} + M(Rq)^{-1/(l+1)} N^{1/(k+1)}) \\ &\ll (N^{1/2} R^{1/2} Q^{3/2} + MN^{1+1/(l+1)} Q^{l/(l+1)} R^{-1/(l+1)}) \log^{2B} Y \\ &\ll Y \log^{2B} Y. \end{aligned}$$

Whence Lemma 5.4 follows.

LEMMA 5.5. *Suppose $1/2 < \gamma_1 < 1$, $\delta = \delta(\gamma_1) = \min((3-2\gamma_1)/14, (\gamma_1-1/2)/2l)$. Then the estimate*

$$S(Y; h_1, \dots, h_l, \gamma_1, \dots, \gamma_l) \ll Y^{1-\delta} \log^{5.5} Y. \quad (5.7)$$

Proof. We sketch the proof. Take $u = Y^{2\delta}$ and $v = Y^{1/2}$ in (3.15). We use Lemma 2.8 to estimate S_1 and S_2 for $m \leq Y^{2\delta}$. We use Lemma 5.4 to estimate S_3 and S_2 for $m > Y^{2\delta}$.

PROPOSITION 5.1. *Let $\delta = \min(\gamma_1/(4l-2), 1/(4l+6))$. Then we have*

$$S(Y; h_1, \dots, h_l, \gamma_1, \dots, \gamma_l) \ll Y^{1-\delta} \log^{5.5} Y. \quad (5.8)$$

Proof. This estimate follows from Lemma 5.3 for $\gamma_1 \geq (2l+3/2)/(2l+3)$ and from Lemma 5.5 for $\gamma_1 \leq (2l+3/2)/(2l+3)$.

6. PROOF OF THEOREM 3

Following the proof of Theorem 1, we only need to estimate $U(h)$ for fixed $h = (h_1, \dots, h_k) \neq (0, \dots, 0)$. We take $H = Y^{\delta_3}$, where δ_3 is defined in Section 1.

For a fixed $h = (h_1, \dots, h_k) \neq (0, \dots, 0)$, let $n_0(h)$ denote the number of h_j such that $h_j \neq 0$, and let d be the first integer j with $h_j \neq 0$. If $n_0(h) \geq 2$, then by Proposition 5.1 we have

$$\begin{aligned} U(h) &\ll Y^{1 - \min(1/(4n_0(h) + 6), \alpha_d/(4n_0(h) - 2))} \log^{5.5} Y \\ &\ll Y^{1 - \min(1/(4k + 6), \alpha_k/(4k - 2))} \log^{5.5} Y. \end{aligned}$$

Now suppose $n_0(h) = 1$. If $\alpha_d \geq 340/531$, then by Proposition 3.1* we have

$$U(h) \ll Y^{1 - 40/407} \log^{11.5} Y \ll Y^{1 - \delta_3} \log^{5.5} Y.$$

If $\alpha_d < 340/531$, then by Proposition 3.3 we have

$$\begin{aligned} U(h) &\ll Y^{1 - \min((1 - \alpha_d)/3, 1/6, \alpha_d/2)} \log^{4.5} Y \\ &\ll Y^{1 - \delta_3} \log^{5.5} Y. \end{aligned}$$

This completes the proof of Theorem 3.

REFERENCES

1. R. C. Baker and G. Kolesnik, On the distribution of p^α modulo one, *J. Reine Angew. Math.* **356** (1985), 174–193.
2. A. Balog, On the fractional part of p^θ , *Arch. Math.* **40** (1983), 434–440.
3. A. Balog, On the distribution of $p^\theta \pmod{1}$, *Acta. Math. Hungar.* **45**, Nos. 1–2 (1985), 179–199.
4. E. Bombieri and H. Iwaniec, On the order of $\zeta(\frac{1}{2} + it)$, *Ann. Scuola Norm. Sup. Pisa* **13**, No. 3 (1986), 449–472.
5. E. Fouvry and H. Iwaniec, Exponential sums for monomials, *J. Number Theory* **33** (1989), 311–333.
6. S. W. Graham and G. Kolesnik, “Van Der Corput’s Method of Exponential Sums,” Cambridge Univ. Press, Cambridge, UK, 1991.
7. S. A. Gritsenko, On a problem of I. M. Vinogradov, *Mat. Zametki* **39** (1986), 625–650. [In Russian]
8. G. Harman, On the distribution of \sqrt{p} modulo one, *Mathematika* **30** (1983), 104–110.
9. D. R. Heath-Brown, The Pjateckskil–Sapiro prime number theory, *J. Number Theory* **16** (1983), 242–266.
10. R. M. Kaufman, The distribution of \sqrt{p} , *Mat. Zametki* **26** (1979), 497–504.
11. J. Schoissengeier, The connection between the zeros of the ζ -function and sequences $(g(p))$, p prime mod 1, *Mh. Math.* **87** (1979), 21–52.
12. J. Schoissengeier, Eine neue Diskrepanz für gewisse Primzahlfolgen, in “Sitzungsber. Math. Natur. Kl.,” Abt. II, 187. Bd., 4. bis 7. Heft, 1978.
13. D. I. Tolev, On the simultaneous distribution of the fractional parts of different powers of primes numbers, *J. Number Theory* **37** (1991), 298–306.

14. R. C. Vaughan, An elementary method in prime number theory, *Acta Arithmetica* **37** (1980), 111–115.
15. I. M. Vinogradov, “Special Variants of the Method of Trigonometric Sums,” Izda. Nauka, Moskow, 1976. [In Russian]
16. W. Zhai, On the k -dimensional Piatetski–Shapiro prime number theorem, *Sci. in China (Ser. A)* **29** (1999), 797–806.